#### §520.1326b

use. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[45 FR 10759, Feb. 19, 1980, as amended at 46 FR 52330, Oct. 27, 1981. Redesignated at 51 FR 13212, Apr. 18, 1986, as amended at 62 FR 61625. Nov. 19. 19971

# § 520.1326b Mebendazole and trichlorfon paste.

- (a) *Specifications*. Each gram of paste contains 100 milligrams of mebendazole and 454 milligrams of trichlorfon.
- (b) Sponsor. See No. 000061 in §510.600(c) of this chapter.
- (c) Conditions of use—(1) Amount. 8.8 milligrams of mebendazole and 40 milligrams of trichlorfon per kilogram of body weight.
- (2) Indications for use. It is used in horses for treatment of infections of bots (Gastrophilus intestinalis and G. nasalis), large roundworms (Parascaris equorum), large strongyles (Strongylus edentatus, S. equinus, S. vulgaris), small strongyles, and pinworms (Oxyuris equi).
- (3) Limitations. Do not administer more than once every 30 days. Do not treat sick or debilitated animals, foals under 4 months of age, or mares in the last month of pregnancy. Trichlorfon is a cholinesterase inhibitor. Do not administer simultaneously or within a few days before or after treatment with, or exposure to, cholinesterase-inhibiting drugs, pesticides, or chemicals. Do not administer intravenous anesthetics, especially muscle relaxants, concurrently. Not for use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[51 FR 13212, Apr. 18, 1986, as amended at 62 FR 61625, Nov. 19, 1997]

### § 520.1330 Meclofenamic acid granules.

- (a) Chemical name. N-(2,6-Dichlorom-tolyl) anthranilic acid.
- (b) Specifications. The drug is in granular form containing 5 percent meclofenamic acid.
- (c) Sponsor. See No. 000856 in §510.600(c) of this chapter.
- (d) Conditions of use. (1) The drug is used in horses for the treatment of acute or chronic inflammatory diseases involving the musculoskeletal system.

- (2) It is administered orally at a dosage of 1 milligram per pound of body weight (1 gram per 1,000 pounds) once daily for 5 to 7 days by addition to the daily grain ration.
- (3) Treatment beyond the initial 5- to 7-day period may be indicated. A maintenance dosage level should be individualized for each animal.
- (4) This drug should not be administered to horses with active gastro-intestinal, hepatic, or renal disease.
- (5) Not for use in horses intended for food.
- (6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 5632, Feb. 9, 1976, as amended at 53 FR 23390, June 22, 1988]

## §520.1331 Meclofenamic acid tablets.

- (a) Specifications. Each tablet contains either 10 or 20 milligrams of meclofenamic acid.
- (b) Sponsor. See No. 000856 in \$510.600(c) of this chapter.
- (c) Conditions of use in dogs—(1) Amount. 1.1 milligrams per kilogram (0.5 milligram per pound) daily for 5 to 7 days.
- (2) *Indications for use*. For the relief of signs and symptoms of chronic inflammatory disease involving the musculoskeletal system.
- (3) Limitations. For oral use only. Should not be administered to animals with congestive heart failure or active gastrointestinal, hepatic, or renal disease. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[50 FR 43385, Oct. 25, 1985, as amended at 53 FR 23390, June 22, 1988]

### §520.1341 Megestrol acetate tablets.

- (a) Specifications. Each tablet contains 5 or 20 milligrams of megestrol acetate.
- (b) Sponsor. No. 000061 in \$510.600(c) of this chapter.
- (c) Conditions of use. (1) The drug is used in female dogs for the postponement of estrus and the alleviation of false pregnancy.
- (2) It is administered orally, intact, or crushed and mixed with food as follows: